



Biological Changes in Human Populations with Agriculture

Author(s): Clark Spencer Larsen

Source: *Annual Review of Anthropology*, Vol. 24 (1995), pp. 185-213

Published by: Annual Reviews

Stable URL: <http://www.jstor.org/stable/2155935>

Accessed: 20/01/2010 15:55

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=annrevs>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



Annual Reviews is collaborating with JSTOR to digitize, preserve and extend access to *Annual Review of Anthropology*.

<http://www.jstor.org>

BIOLOGICAL CHANGES IN HUMAN POPULATIONS WITH AGRICULTURE

Clark Spencer Larsen

Department of Anthropology and Research Laboratories of Anthropology, The University of North Carolina, Chapel Hill, NC 27599-3120

KEY WORDS: skeletal biology, nutrition, technology, demography, disease, activity

ABSTRACT

Agriculture has long been regarded as an improvement in the human condition: Once *Homo sapiens* made the transition from foraging to farming in the Neolithic, health and nutrition improved, longevity increased, and work load declined. Recent study of archaeological human remains worldwide by biological anthropologists has shown this characterization of the shift from hunting and gathering to agriculture to be incorrect. Contrary to earlier models, the adoption of agriculture involved an overall decline in oral and general health. This decline is indicated by elevated prevalence of various skeletal and dental pathological conditions and alterations in skeletal and dental growth patterns in prehistoric farmers compared with foragers. In addition, changes in food composition and preparation technology contributed to craniofacial and dental alterations, and activity levels and mobility decline resulted in a general decrease in skeletal robusticity. These findings indicate that the shift from food collection to food production occasioned significant and widespread biological changes in human populations during the last 10,000 years.

INTRODUCTION

For much of Western history, the popular and scholarly perception of agriculture has been that once it was acquired, *Homo sapiens* had it made—life improved dramatically from a state of incessant work to abundant leisure, from

deprivation to plenty, and from sickness to health and increased longevity. Moreover, the burden of constantly moving about the landscape in search of food was replaced by the security of settlement in permanent towns and cities from which the many advantages of civilization could be enjoyed (e.g. 24). Beginning in the late 1960s, a body of data began to emerge suggesting that this characterization of the shift from foraging to farming is inaccurate, and many hunter-gatherers may not have had it all that bad, especially with regard to work load or health (142, 241). Some researchers began to speculate that the adoption of an agricultural lifeway may not have been a positive development, at least with regard to its health costs (52).

Until recently, the literature lacked data that could be used to document and interpret biological changes in human populations with the transition from food collection to food production. Over the past decade or so, biological anthropologists (e.g. 55) have marshalled a wealth of data pertaining to human biological change in relation to the shift from foraging to farming. This record indicates that commonly held assumptions about the benefits of the agricultural transition for humankind are in need of reassessment (8, 53, 54).

This review relies on diachronic and other comparative studies of human skeletal remains recovered from archaeological sites in order to summarize and interpret biological changes with the shift from foraging to farming. Like other types of archaeological data sets, there are biases—such as cultural practices, differential preservation, excavation strategies, and selective curation—that potentially confound attempts to accurately reconstruct and interpret the past (211, 233, 256, 263). When these biases are evaluated carefully, however, human remains provide a valuable retrospective picture of health and behavior in the past that is not available from any other source of information (136).

For this review, I define agriculture broadly to include any type of plant cultivation practiced for dietary purposes (73). Independent centers of domestication occurred on all inhabitable continents (except Australia) within the last 10,000 years (93), first in the Old World and followed in the New World several thousand years later (74). Major crops included barley and wheat in the Near East; millet, sorghum, yams, and dates in Africa; millet and rice in northern China; rice, sugarcane, taro, and yams in Southeast Asia; maize, beans, and squash in Central America; and potatoes, sweet potatoes, and manioc in South America. Use of domesticated plants spread to adjacent regions (e.g. maize into the American Southwest and eastern North America and wheat and barley into Europe) at varying rates and involving localized circumstances (50, 58, 79, 93, 212).

Consideration of these centers and associated regions of domestication reveals a number of common themes that have implications for human biology, especially those linked to changes in health and behavior. This review is

organized around the following themes: (a) composition and nutritional quality of food, (b) food processing technology and mastication, (c) population growth, (d) population distribution and health, and (e) work load and activity patterns. These themes are not mutually exclusive; health status, for example, was highly influenced by a number of factors, including nutrition, population distribution, and behavior. In addition, although biological changes have been identified with agricultural shifts worldwide, there are some notable exceptions, even within the same region. From 250 B.C. to A.D. 200 in the Eastern Woodlands of North America, use of cultivated seed-bearing plants (e.g. chenopod, sunflower, and cucurbit) coincided with social and economic transformations (222). Later, between A.D. 800 and 1100, the emphasis shifted to a single plant, maize (221). The earlier economic transition to seed plants appears to have had some influence on human biology, such as bone structural morphology and dental health (31, 200; see also 75). However, based on studies of numerous skeletal samples, maize agriculture had a far more profound impact on human populations than the earlier dietary regime, at least with respect to health status, demographic history, and activity (see below).

Some domesticated plants (e.g. maize, millet) consumed by past groups leave detectable chemical and isotopic signatures in human remains. The documentation of changes in these signatures has greatly increased the precision of the timing of the shift to agriculture in various human populations. The primary focus of this discussion is on health consequences and anatomical changes and not on the chronology of subsistence change. Therefore, bone chemistry and dietary reconstruction are not addressed here (see 125, 208, 214).

FOOD COMPOSITION AND NUTRITIONAL QUALITY

Dental Health

One of the most striking changes to occur with the adoption of agriculture is a decline in dental health, resulting largely from an increasing emphasis on carbohydrates in the diet. These changes are best illustrated in prevalence of dental caries and antemortem tooth loss.

DENTAL CARIES Dental caries is a disease process involving focal demineralization of dental hard tissues by organic acids produced by bacterial fermentation of dietary carbohydrates, especially sugars (176). Numerous workers have detailed increases in carious lesions (cavities) in agriculturalists compared to hunter-gatherers in North America, South America, Europe, Asia, and Africa (reviews in 55, 141, 165, 187, 242). From a sample of populations drawn

globally, Turner (242) determined average frequencies of teeth affected (incisors + canines + premolars + molars): foragers, 1.7%; mixed (agriculture + foraging), 4.8%; and agricultural, 8.6%. Frequencies within each of these categories are highly variable, however. Few studies comparing caries prevalence for different types of cultigens (e.g. maize vs wheat) have been made. Based on caries prevalence data, Lubell and associates (144) observed that European Neolithic domesticates were either less cariogenic or had less dietary importance than maize in North America.

The composition of food and the manner of its preparation have been linked to higher caries prevalence (192). Many agricultural societies prepare plant foods by boiling them, usually into a soft, gruel-like consistency. Consumption of these foods promotes the growth of bacterial colonies in fissures and grooves of premolars and molars and other areas of tooth crowns that are not cleansed mechanically by the action of saliva or mastication.

Some researchers have noted apparent negative correlations between caries prevalence and degree of occlusal wear in human populations with cariogenic diets (150, 165, 192). However, Meiklejohn and coworkers (155, 159) indicate that although a high-wear environment may be a cariostatic factor, dental caries and wear are independent variables and a relationship between them should not be generalized. For instance, various populations with rapid wear-rates also exhibit significant prevalence of dental caries resulting from consumption of highly cariogenic foods (94, 159).

Dental caries clearly increases with agriculture, but appreciable frequencies of carious lesions are not unique to human groups relying on starchy domesticated plants. For example, carious lesions are common in hunter-gatherer dentitions from the lower Pecos region of the Chihuahuan Desert of Texas, which is likely the result of reliance on high-carbohydrate plants with a sticky texture (e.g. succulent fibers, pecans, prickly-pear fruits) (94, 228). High caries prevalence in preagricultural Mesolithic series from Sicily and Portugal compared to other contemporaneous European dental samples (e.g. 156) points to the consumption of cariogenic nonagricultural foods (e.g. honey) or sweet, sticky fruits (e.g. dates or figs) (19, 70, 144, 155). Comparison of Mesolithic and Neolithic dentitions from Portugal indicates that the latter group did not have higher caries rates than their foraging-fishing predecessors (144, 155). Lukacs (146) has reported a high prevalence of caries in the Langhnaj Mesolithic sample from India, which he suggests may reflect inclusion of agricultural foods obtained through trade or exchange with nearby agriculturalists. Frequency of carious lesions declined from high to low levels on Santa Rosa Island, California, when groups shifted from reliance on nondomesticated plants (e.g. roots and tubers) to a marine-based diet (254). Levels of natural fluoride in ground water also appear to be associated with variation in dental

decay; when diets are cariogenic, high fluoride prevents caries and low fluoride promotes it (148, 174, 219).

Furthermore, the increase in dental caries with agriculture was greater in women than in men in most regions (133, 141, 147, 255), which indicates widespread gender-based differences in preparation and consumption of food. Some evidence indicates that a combined high-carbohydrate and low-protein diet in females may have predisposed their teeth to more decay than males (132).

TOOTH LOSS Antemortem tooth loss is caused by a variety of factors. For example, periodontitis (periodontal disease) causes degeneration of the alveolar bone and other tissues anchoring the teeth in the jaws. As the resorption process progresses, the bony support is reduced, resulting in exfoliation of teeth (51, 96, 98). As with dental caries, consumption of soft carbohydrate foods promotes periodontal disease. Many studies documenting caries increase with the transition to farming or agricultural intensification also report elevated levels of tooth loss, suggesting a strong relationship between caries and tooth loss (e.g. 9, 11, 56, 91, 123, 147, 227; but see 48, 106, 157). Rapid or high wear-rates, pulpal exposure, and excessive buildup of plaque also contribute to antemortem tooth loss (51, 94).

Growth and Development

The shift from foraging to farming appears to have resulted in a decline in nutritional quality for many human groups. In this regard, agricultural populations consumed a narrower range of foods than did hunter-gatherers; this narrowing of the diet involved reduced availability of animal protein coupled with dependence on a limited number of domesticated plants. Archaeological documentation of prehistoric farming diets and observation of living peasant agrarian populations indicate that agricultural diets tend to be dominated by one or a few plants, such as rice in Asia, wheat in temperate Asia and Europe, millet or sorghum in Africa, and maize in the New World.

The nutritional value of some of these plants is marginal or poor. Maize, for example, is deficient in the essential amino acids lysine, isoleucine, and tryptophan (67). Niacin (vitamin B₃) in maize is chemically bound, which reduces the bioavailability of this nutrient to the consumer. Moreover, iron absorption in maize-based diets is exceedingly low (10). In many regions of the Old World, milled grains such as millet and wheat contain very little iron. In some settings (e.g. the Nile Valley), weaned children ate cereal grains containing virtually no iron (45). Rice is deficient in protein, even in the unmilled form. Moreover, the low availability of protein in rice inhibits vitamin A activity, even if the vitamin is provided through other food sources (261).

Many societies have developed means of improving the nutritional content of these so-called superfoods, such as alkali-processing of maize (120, 230). Nevertheless, the effects of over-reliance on these plants are profound, which is indicated by the anthropometric histories of individual populations, including secular changes in growth and adult body size (66).

GROWTH RATES Growth rates in children provide a highly reliable indicator of nutritional status (88). Studies of living populations experiencing nutritional deficiencies show that children are short for their age in comparison to populations with adequate diets (18, 66, 112). These findings underscore the great sensitivity of childhood growth to environmental factors, which affect all body tissues, including the skeleton (18).

Analysis of juvenile long bones from prehistoric North America reveals evidence of growth retardation in agricultural and mixed-subsistence economies (210). In children less than six years of age in the lower Illinois Valley, matching of femur length to dental age reveals that growth is impeded in incipient agriculturalists as compared to earlier foragers (56). Interestingly, individuals who are short for their age tend to have a higher frequency of physiological stress indicators [e.g. *cribra orbitalia* (lesions in the eye orbits) and tooth enamel defects] than do individuals who are tall for their age, lending further support to nutritional deficiency as a factor contributing to growth retardation (56). Similarly, in the late prehistoric Dickson Mounds populations from west-central Illinois, growth of long bones (humerus, tibia, femur) from birth to adulthood declined appreciably (especially for two- to five-year-olds) during the period of intensive agriculture relative to earlier periods of less intensive agriculture (83). Circumferences of long bones express an equivalent pattern of retarded growth, although the greatest discrepancy between the later intensive agriculturalists and the earlier population at Dickson Mounds was between ages 10 and 15 (83).

Mortality bias in cemetery samples—especially that involving the inclusion of a relatively greater number of physiologically stressed children—might present a skewed picture of juvenile body size and growth retardation. One would expect to find a higher proportion of sick (and shorter) children in cemetery samples. Saunders & Hoppa (211), however, demonstrate that although mortality bias exists in archaeological juvenile remains, the effects are small and are far outweighed by errors introduced by other factors (e.g. aging methodology, sample size, and preservation status).

STATURE Substantial evidence drawn from the study of living populations reveals the strong relationship between growth retardation and attainment of adult body size, including terminal height (e.g. 72). That is, growth-retarded children should be short-statured adults. Abundant evidence shows that stature

is linked with environmental factors. Comparison of successive periods (e.g. by decade) in living or historically documented populations indicates that during periods of increased dietary stress and food shortages, stature of children and adults reduces (68, 231, 232, 265); with improved nutrition, stature increases (66).

Several geographic settings show stature declines with agriculture adoption or intensification in prehistoric societies (7, 83, 123, 132, 135, 157, 178, 188). In contrast, other regions show increases or no change in stature (48, 56, 198, 246, 248). Thus, there is evidence for stature reduction with agriculture in selected settings, but this is not a universal pattern. The ambiguous picture of stature change in comparisons between prehistoric populations adopting agriculture may involve differences in access to critical dietary resources (e.g. protein) during years of growth that are especially sensitive to environment (e.g. the period of catch-up growth in adolescence). Thus, in some populations, nutritionally deprived children may have had better diets during adolescence, resulting in growth rebounds and, hence, full growth and normal attainment of stature as adults (e.g. 231).

CORTICAL BONE THICKNESS A strong relationship has been demonstrated between nutritional quality and bone mass or thickness in tubular bones (e.g. femur, metacarpals). Losses in cortical bone thickness in nutritionally stressed living populations have resulted from reduced growth as well as from increased endosteal (interior bone surface) resorption (76, 78, 99). In archaeological remains, temporal changes in or characteristics of cortical thickness have been related to nutritional status (56, 110, 111, 183). That is, relatively thin cortical bone frequently is interpreted as reflecting poor nutrition.

Nutrition is linked closely to skeletal maintenance. However, the mechanical environment is also an important consideration in interpreting cortical bone thickness and distribution. The external surface (subperiosteum) and internal surface (endosteum) of cortical bone continue to expand or become more outwardly distributed throughout adulthood (76). This expansion is influenced strongly by the level of mechanical or functional demand that occurs during one's lifetime. Cortical thickness frequently is calculated as percent cortical area (PCCA) or percent cortical thickness (PCCT), which are simple measures of the amount of cortical bone relative to the subperiosteal area or overall breadth of the bone (e.g. 76). Declines in these values in archaeological remains may represent decreases in nutritional status (e.g. 56). However, both the subperiosteum and endosteum may be expanded, resulting in considerably reduced PCCA or PCCT (201). In these circumstances, although there is less bone tissue relative to outer dimensions, the outward redistribution of bone provides greater mechanical strength (see below). Thus, in interpreting pat-

terns of cortical thickness, both systemic (nutritional) and mechanical (structural) factors need to be evaluated carefully.

BONE HISTOLOGY Like any other living tissue, bone continually renews itself. The renewal process is unique in bone and involves resorption followed by deposition—specialized cells called osteoclasts remove bone, and osteoblasts replace it. The histologic structures, osteons or Haversian systems, associated with this process can be measured precisely, revealing important information on health status and nutritional quality in earlier societies (153, 235). Stout (234) has compared bone-remodeling rates of prehistoric hunter-gatherers and maize agriculturalists in North America (Gibson, Ray, and Ledders sites, Illinois) and South America (Paloma, Peru). This histomorphometric analysis documents greater remodeling rates in maize agriculturalists (Ledders) than in hunter-gatherers, which Stout suggests may reflect the effects of overproduction of parathyroid hormone resulting from the low calcium/high phosphorus ratios characteristic of high maize diets.

To compensate for bone losses in aging adults (especially after age 40), structural adaptations may occur involving more outward distribution or expansion of bone tissue (202, 205). Greater remodeling rates may supplement these changes (39). Thus, remodeling differences may reflect mechanical responses that are not necessarily tied to nutritional factors.

TOOTH SIZE Tooth size appears to be under greater genetic control than is bone size (124). The pattern of reduction in human tooth-size over the course of hominid evolution (22, 23, 41, 42, 69, 124, 145) suggests that this is largely evolutionary (genetic) change, although the mechanism remains elusive (124). This trend of tooth-size reduction has continued in the Holocene (but see 115, 216) and has been related alternatively to dietary change and the adoption of agriculture, use of pottery and cooking, increasing sedentism and population density, migration and gene flow, craniofacial or body-size variation, dental disease, or other factors (e.g. 22, 23, 41, 42, 63, 89, 132, 145, 151, 224).

The presence of small teeth in physiologically stressed individuals in living populations suggests that environment (e.g. nutritional or maternal health status) also influences tooth size (77, 239). Therefore, like body size, tooth size should provide a measure of deviation from genetic growth potential. However, unlike body size, teeth are not subject to catch-up growth or rebound during the adolescent growth spurt. Prehistoric maize agriculturalists on the southeastern US Atlantic coast have smaller teeth than did their foraging predecessors (134). Because deciduous tooth crowns are formed largely in utero, smaller teeth in the agricultural population may reflect a reduction in maternal health status and placental environment as a result of consumption of

nutritionally poor food (maize). Reduction in permanent-tooth size has also been observed in this setting (132) and elsewhere (e.g. 63, 105, 160) with the shift to agriculture. Given the narrow temporal window of tooth-size reduction in recent past populations (less than several hundred years), and especially when viewed in the context of reduced health and dietary quality, these changes may reflect, at least in part, decline in nutritional status. This does not mean, however, that tooth-size reduction can be explained fully by nonevolutionary factors. Calcagno (41), for example, has documented a relatively greater reduction in posterior-tooth size than anterior-tooth size in Nubian populations. He has tied this change to a selective advantage for smaller posterior teeth in caries-prone agriculturalists.

The years of growth and development of the dentition (4 months in utero to age 12) involve a period of elevated stress for human populations (220). Therefore, individuals experiencing relatively greater stress during this period should have smaller tooth crowns. Comparison of crown dimensions of pre-adult and adult age-at-death cohorts shows that preadults had smaller permanent teeth than did adults (90, 220). These studies document the failure of teeth to reach their genetic size potential in nutritionally stressful settings. Moreover, they imply that individuals with small teeth had a reduced lifespan, a finding that is consistent with other hard-tissue investigations (e.g. 49, 56, 81). It is unlikely, however, that small teeth led to reduced longevity. Rather, size reduction is symptomatic of environmental factors (e.g. undernutrition) that contributed to smaller teeth and perhaps to premature death.

DENTAL ENAMEL DEFECTS Various kinds of dental enamel defects are represented abundantly in past populations (82, 86, 87). The most frequently observed defects are hypoplasias, which are deficiencies of enamel expressed as circumferential lines, grooves, or pits resulting from the death or reduced function of enamel-producing cells (ameloblasts) and failure to form enamel matrix (86). A suite of causes of enamel defects have been identified, including various infectious diseases and nutritional deficiencies (86, 87). Studies of living populations demonstrate a strong link with nutritional status (84).

Comparisons between foragers and farmers generally reveal the latter to have more hypoplasias, which is interpreted to represent an increase in physiological perturbation (48, 56, 83, 123, 188, 195, 198, 247, 248; but see 106, 266). One investigation has documented a decrease in individuals affected by hypoplasias, but agriculturalists have wider hypoplasias, suggesting that there was a reduction in number of stress events, which were either longer or more severe following the adoption of maize (113, 138; see also 17). Other studies of agricultural populations, both prehistoric and contemporary, reveal elevated frequencies of hypoplasias in nutritionally compromised individuals with low socioeconomic status (85, 259). Goodman and coworkers (83) reported that

stress in agricultural children peaked earlier than in hunter-gatherer children from the same archaeological locality in the American Midwest. Thus, physiological stress is higher and socially and demographically patterned in this agricultural population.

Stress-related histological defects of tooth enamel (e.g. Wilson bands and other microdefects) accord well with hypoplasia prevalence in comparisons between hunter-gatherer and agricultural populations (86; but see 61, 199). Rose and colleagues have found a fourfold increase in enamel microdefects in intensive agriculturalists in comparison to incipient agriculturalists in Illinois; they link this increase to a decline in nutrition and an increase in susceptibility to infection (85, 128, 197, 199). Although many studies of physiological stress in prehistoric agriculturalists identify nutrition as a primary cause, most acknowledge the important synergy between poor diet and infection: Undernourished individuals are more prone to infection, and infection detrimentally affects nutritional status (218).

FOOD PROCESSING AND MASTICATION

Dental Attrition and Trauma

Mastication involves preparation of food with the incisors and canines followed by reduction of food with the premolars and molars. As a result, the teeth wear as they come into contact with food and with each other. The wear rates are highly influenced by the consistency and texture of food, which is determined either by the characteristics of the food itself (e.g. presence of phytoliths or cellulose in plants) or by the manner of its preparation. Plants are processed in a number of different ways (230), some of which may involve the introduction of abrasive elements that promote tooth wear (e.g. use of grinding stones for making flour from cereal grains). Wear on tooth surfaces is documented both visually (macrowear) and microscopically (microwear).

MACROWEAR Wear on the chewing surfaces of teeth varies widely between human populations, owing to localized behavioral characteristics, cultural practices, age, sex, and diet (173, 185, 253). Because many Western populations consume soft, highly processed foods that include little extraneous material, tooth wear proceeds exceedingly slowly from birth through adulthood. In many nonindustrial societies, however, grit inclusions and minimal processing of food contribute to rapid wear rates (94).

Tooth wear is strongly influenced by age: Older individuals have been exposed to factors resulting in attrition for a longer period of time than have younger individuals. Because of the difficulty of age determination in archaeological human remains, wear rates are quantified imprecisely (97, 217). How-

ever, valuable information can be generated by comparison of age groups within populations (e.g. five-year intervals). Comparisons between earlier hunter-gatherers and agriculturalists or populations undergoing agricultural intensification reveal general declines in degree or severity of tooth wear in a wide range of settings (1, 2, 9, 12, 14, 48, 105, 107, 109, 114, 123, 144, 177, 184, 185, 187, 191, 192, 200, 216, 225, 251). This trend reflects the reduction in consistency, hardness, or abrasiveness of foods consumed by agriculturalists. For many regions, the introduction of ceramic vessels and accompanying food-preparation practices involving extended cooking engendered reduction in food consistency (23, 172). Some settings involving agriculture had an increase in use of grinding stones, thus resulting in relatively more rapid occlusal wear than in preagricultural peoples (e.g. 171, 226).

Types or patterns of wear have also changed in the shift to agriculture. Smith (223) has found a dichotomy of occlusal wear patterns in molars: Hunter-gatherers have evenly distributed flat wear, and agriculturalists have highly angled wear. The change in wear pattern in comparisons between earlier foragers and later farmers was interpreted to reflect a reduction in "toughness" of agricultural diets. Other researchers have confirmed this finding but with some notable exceptions (e.g. 144, 213, 215).

In specific regions additional distinctive wear patterns have been documented, including the presence of cupped wear on occlusal surfaces of molars (144, 184, 223) and incisors (100) of agriculturalists but not hunter-gatherers. Hinton (102) has identified greater interproximal wear (wear on nonocclusal surfaces of adjacent teeth) in prehistoric hunter-gatherers from Tennessee, which he relates to increased magnitude and frequency of masticatory forces in hunter-gatherers (also 187).

MICROWEAR Recent application of microscopic techniques, especially scanning electron microscopy (SEM), to the study of tooth wear has facilitated understanding of dietary change and adoption of agriculture in past populations (171, 238). Comparisons between hunter-gatherers and agriculturalists reveal a decrease in frequency of microwear features (e.g. pits and scratches) on occlusal surfaces (38, 107, 191, 200, 238). Like the macrowear evidence, this finding indicates consumption of softer foods, which results in fewer microscopic features, such as pits, in agriculturalists than in hunter-gatherers.

In contrast, comparison of Mesolithic, Neolithic, and Chalcolithic microwear in dental series from South Asia and Mesolithic and Neolithic dental series from northern Syria shows an increase in frequency of microwear features, especially large pits (171, 184–186). This pattern appears to reflect the use of coarse grinding stones by Neolithic and later agriculturalists in preparing cereal grains, and the shift from consumption of small-grained seeds to

large-grained cereals. The pattern also reflects the shift from roasting meat directly on coals or air-drying to cooking with ceramic vessels. Analysis of later agriculturalists (post-5300 B.C.) from Syria reveals a reversal of this trend. That is, there is a sharp decrease in frequency of microwear features, which appears to be related to the introduction of ceramics and cooking of cereals into soft mushes (172).

DENTAL CHIPPING Some populations consuming abrasive diets display fractures, cracking, and other evidence of damage to tooth crowns. A number of foraging groups consume tough foods containing various abrasives and, as a result, have a relatively high frequency of damaged teeth (19, 244). Milner & Larsen (169) have reviewed reports on dental trauma, but few studies have indicated differences between hunter-gatherers and agriculturalists. Patterson (187), however, observed a marked decline in dental trauma in comparisons between prehistoric foragers and farmers from Ontario. He interpreted this trend as reflecting a reduction in the use of teeth for food processing in the agriculturalists. Comparisons between foraging and farming dentitions in East Asia suggest a decline in dental trauma (243).

Craniofacial Adaptation

CRANIOFACIAL GRACILIZATION A trend in recent human evolution toward a decrease in face size has long been recognized by anthropologists and others. In pre-twentieth-century British populations, for example, Keith (32:198) observed "maxillary shrinkage" and facial reduction in recent populations compared to those of earlier periods. He attributed these changes to consumption of "cooked food and soft cereals replacing tough meats and imperfectly ground corns." Recent studies have reported a consistent pattern of reduction in craniofacial robusticity and/or vault shape in populations that underwent the transition from foraging to farming, both in the Old World (41, 46, 47, 62, 64, 114, 227, 237, 264) and in the New World (2, 20, 101, 132, 136, 177). To varying degrees, changes are similar to those documented in Nile Valley crania by Carlson and colleagues (44, 46, 47, 103, 104, 249, 250), who argue that craniofacial alterations (i.e. shorter and rounder cranial vaults, smaller and more posteriorly placed faces, general reduction in size and robusticity of faces and jaws) resulted from "progressive alterations in maxillomandibular growth in response to developmental variation in the size and position of the muscles of mastication" with a shift to the consumption of soft, agricultural foods (47:574).

MALOCCLUSION Other orofacial changes accompanying shifts from hard-textured to soft-textured foods include tooth crowding and malocclusion. Although relatively little research has been devoted to explaining underlying

causes of malocclusion, a great deal of information exists on occlusal variation in humans, past and present (see 57). In general, nonindustrial groups masticating and ingesting hard foods possess "edge-to-edge" bite, whereby the upper teeth come into greater occlusal contact with lower teeth. In contrast, industrial or other groups consuming soft foods have high frequencies of occlusal abnormalities and crowding (e.g. overjet, overbite, crossbite, tooth impaction, tooth rotation) (57). Mechanisms for occlusal abnormalities are controversial, but experimental studies involving controlled feeding of soft- and hard-textured foods to laboratory animals indicate the primacy of food consistency in interpreting prevalence of malocclusion: Consumption of predominantly soft foods promotes the development of occlusal abnormalities (57).

A large body of data exists on occlusal abnormalities in living human populations, including nonindustrial societies (57). Early contact period Eskimo and Aleut populations pursuing traditional dietary practices had a low prevalence of malocclusion; when food consumption practices shifted to softer westernized diets, dental crowding and occlusal variations increased (175, 262). Diachronic changes in archaeological settings have been reported only minimally. However, evidence from the eastern Mediterranean (3), South and East Asia (92, 123, 149, 237), and the American Southeast (177) indicates that the shift to agriculture or more intensified agriculture was accompanied by an increase in dental crowding and malocclusion.

POPULATION GROWTH

One of the most significant changes in human evolution is the dramatic increase in population size during the Holocene, especially in areas adopting agriculture (8, 53, 95). Various causes have been proposed to explain this growth, but evidence suggests that increase in fertility and birthrate plays a central role. However, this is a point of contention among anthropologists and demographers (53). A fertility-based argument for population increase is consistent with the generally greater fertility seen in agriculturalists as opposed to nonagriculturalists in ethnographic settings (15, 16), albeit with a high degree of heterogeneity (43).

Paradoxically, skeletal series of agriculturalists show lower mean age-at-death than do hunter-gatherers, which generally has been interpreted to reflect an increase in mortality and declining life expectancy with the shift to agriculture (83, 123, 126, 135, 258). Reevaluation of demographic profiles in skeletal series suggests, however, that mean age-at-death is related to fertility and birthrate, and not mortality (209; see also 36, 116, 140, 158, 168, 172). These studies suggest that human populations experiencing growth will have a greater number of younger individuals, resulting in a larger proportion of

juvenile skeletons relative to adult skeletons. Thus, population growth appears to influence the distribution of age-at-death in cemetery assemblages.

In the lower Illinois River region, birthrate (which is inversely proportional to the ratio of number of deaths over age 30 to number of deaths over age 5) shows an increase prior to, and especially during, the adoption and intensification of maize agriculture in late prehistory (36). Mortality shows no change over the same period. Although various factors are likely involved, the shift to a high-starch diet coupled with improved ceramic technology and cooking techniques would have facilitated earlier weaning and decreased birth interval because of the availability of soft, easily digestible foods (36; cf 172).

Other evidence suggests that the shift to agriculture was accompanied by a general increase in birthrate. In the comparison of hunter-gatherer and agricultural skeletal samples from throughout North America, Buikstra & Konigsberg (35) found that foragers have mortality curves that rise rapidly with increasing age; agriculturalists have much flatter mortality curves. This pattern indicates a younger sample (and thus greater fertility) among agriculturalists than hunter-gatherers.

Although usually difficult to identify, factors contributing to mortality can be recognized in osteological samples. As discussed above, various indicators of morbidity are well represented in skeletal series (e.g. infectious disease). Moreover, traumatic death resulting from warfare and conflict have been documented in prehistoric osteological samples worldwide, with some of the best evidence coming from North America (e.g. 130, 167, 260). This evidence indicates that a number of late prehistoric agricultural societies were involved in conflict situations resulting in an increased risk of early death. Reports on skeletal trauma arising from malevolent activities suggest that there was an increase in frequency of deaths arising from conflict and warfare among late prehistoric agriculturalists compared with earlier groups (167). The causes of conflict and warfare in these late prehistoric settings are difficult to determine, but competition for increasingly limited resources (e.g. arable or otherwise productive lands) may have been a factor (167).

SETTLEMENT AND POPULATION DISTRIBUTION

Infectious Disease

Reduced population mobility and increased aggregation provide conditions that promote the spread and maintenance of infectious and parasitic diseases and the increase in pathogen load in humans. That is, closer, more crowded living conditions facilitate greater physical contact between members of a settlement, and permanent occupation can result in decreased sanitation and hygiene. Diachronic comparisons of prevalence of nonspecific infections (or

generalized skeletal inflammations called periosteal reactions or periostitis) and some specific infectious diseases (e.g. tuberculosis, treponematoses) indicate that more densely settled agricultural societies were more prone to infection than were earlier groups (7, 37, 48, 56, 83, 128, 132, 135, 137, 157, 188, 198, 200, 246, 248; but see 106, 180). These changes did not result directly from shift in diet, but rather from increased sedentism and aggregation of population and general deterioration of living conditions. This conclusion is corroborated by the presence of similar patterns of increased infection rates in nonagricultural groups that became less mobile. For example, in the Santa Barbara Channel Islands of coastal southern California, increasing population density and declining mobility were accompanied by an elevated prevalence of periostitis (129, 131). Sedentism, however, does not explain fully the general pattern of very high infection rates observed in some prehistoric agriculturalists (55, 65, 166, 167). In this regard, the synergy between infection and other stressors—e.g. poor nutrition, warfare, anemia, social disruption—is key to understanding patterns of remarkably poor health in some late prehistoric agricultural groups.

Anemia

Many archaeological skeletal series have appreciable frequencies of sieve-like cranial lesions in the eye orbits (called *cribra orbitalia*) and flat vault bones (called *porotic hyperostosis*). These lesions typically are accompanied by expanded blood-forming marrow, indicating that the condition represents a response to various types of anemia and increased production of red blood cells (236; cf 182). In archaeological remains, *cribra orbitalia* and *porotic hyperostosis* have been attributed mostly to iron deficiency anemia caused by consumption of iron-poor foods, parasitism, infantile diarrhea, and other chronic stressors that influence iron metabolism (80, 83, 236).

Studies have documented higher prevalence of either *cribra orbitalia* or *porotic hyperostosis* in sedentary farmers compared with mobile foragers (4, 5, 7, 45, 48, 56, 83, 123, 127, 180, 188, 195, 198). There are exceptions to this pattern, however, pointing to factors influencing iron bioavailability that do not involve subsistence directly. Foragers from North America's Pacific coast, for example, have a relatively high prevalence (59, 60, 131, 252) and, conversely, some agriculturalists show very low prevalence of *cribra orbitalia* and *porotic hyperostosis* (140, 193). These situations indicate that a range of nondietary factors are important in the etiology of iron deficiency anemia in past populations (45, 80, 161, 196).

WORK LOAD AND ACTIVITY

There has been an ongoing discussion in anthropology about the demands of labor and other physical behaviors in foraging vs farming lifestyles (52, 53). Study of skeletal series from hunter-gatherers and later agriculturalists has provided abundant information on the biological concomitants of activity, especially with respect to degenerative pathology in articular joints, other articular joint modifications arising from specific and generalized types of activity, and structural adaptations of long bones.

Osteoarthritis and Osteophytosis

Osteoarthritis is a disorder involving the mechanical degeneration of skeletal articulations, resulting in buildup of bone along joint margins (lipping), loss of bone on joint surfaces, or a combination of both. Degenerative changes involving the margins of vertebral disks results in lippling or osteophytosis (30). Because degenerative changes in joints result largely from physical demands occurring over the course of an individual's lifetime (108), their prevalence in past populations provides an important perspective on activity in both living and extinct societies. Observations on osteoarthritis and osteophytosis have been made by anthropologists and others since the nineteenth century, including those comparing foragers and farmers (30). Recent studies have documented decreases in prevalence, severity, or both in a number of settings, including the southeastern United States (28, 132, 134), South Asia (123), and Europe (157). Comparisons of skeletal samples from different regions have shown that although the results are highly variable, hunter-gatherers tend to have more osteoarthritis and osteophytosis than do agriculturalists (117–119, 181, 190). On the other hand, some settings show either no difference or greater frequency of these disorders in agriculturalists than in hunter-gatherers (30, 83, 152, 189). In summary, these studies indicate that although a number of populations underwent change in articular degenerative pathology and presumably also in the behaviors that caused it, no definitive pattern relating prevalence and subsistence mode emerges except with regard to specific geographic settings.

Other Activity-Related Articular Modifications

SPONDYLOLYSIS Spondylolysis is a bilateral (or rarely, unilateral) separation of vertebral neural arches in the area between the superior and inferior articular processes (called the pars interarticularis), mostly in lumbar vertebrae. No other primate shares this trait with humans, suggesting that bipedality and associated behaviors play an important role in its etiology (26, 163). In living populations, high frequencies of spondylolysis are found among individuals who participate

in activities that entail heavy mechanical demands on the lower back, suggesting that it is a type of fracture (26, 163).

Eskimos have an unusually high spondylolysis prevalence, which is consistent with their physically demanding lifestyle in traditional settings (e.g. 162). Bridges (26) has documented a higher prevalence in foragers than in later farmers in prehistoric Tennessee. She notes that although an association of specific activities with spondylolysis is not possible, the defect may be related to specific activities or postures rather than to overall activity levels.

SPECIFIC TASKS AND SKELETAL MODIFICATIONS Distinctive patterns of degenerative joint disorders (especially osteoarthritis) have been identified in hunter-gatherers and agriculturalists. In prehistoric populations from the American Southwest, bilateral elbow osteoarthritis is commonplace in some populations, especially in women (164). This pattern reflects physical activities requiring the use of both arms, such as preparation of cereals into flour with grinding stones. In prehistoric agricultural populations in Syria and Ecuador, the ends of the first metatarsals show extensions of articular surfaces, indicating extreme bending of the toes backward, apparently in kneeling postures (170, 245). With regard to the Syrian evidence, tomb art depicts an individual with toes bent backwards while grinding cereals in a stone quern (170).

Other behaviors that may be unique to prehistoric foragers have been reconstructed from size and robusticity of muscle attachment areas in arm bones. Kennedy (122), for example, has found that terminal Pleistocene and Mesolithic hunter-gatherers in South Asia commonly exhibit well-developed supinator muscle attachment areas on the ulna (one of the two bones of the forearm). He suggests that spear throwing was the activity associated with this feature. The lower frequency of the hyperdeveloped supinator attachment site in later agriculturalists suggests that activities involving heavy demands on the lower arm declined appreciably with the adoption of agriculture.

Robusticity and Structural Adaptation

GENERAL ROBUSTICITY PATTERNS The size and structure of cortical bone is highly sensitive to mechanical stimuli (154). Accordingly, human populations experiencing demanding physical regimes have larger and more robust skeletal elements than do sedentary or less-active populations (201, 207). Comparisons between hunter-gatherer and farming populations from archaeological settings indicate a pattern of decrease in dimensions of long-bone shafts and muscle attachment sites, which presumably reflects a decline in physical demand and work load following sedentism (132, 135, 188, 227; but see 21, 25, 27).

Some investigators have compared long-bone cross-sections from hunter-gatherers and agriculturalists. In many cases, the shafts of bones (e.g. femur) of agriculturalists are more circular in cross section (13, 109, 132, 134, 188). Increasing circularity of long-bone shafts is apparently part of a worldwide trend since the late Pleistocene (34, 143, 203). In hunter-gatherers, femur midshafts (mid-thigh region) and femur proximal shafts (hip region) tend to be more compressed from side-to-side and from front-to-back, respectively, than in agriculturalists or sedentary groups (132, 206). This flattening of long bones in foragers has been interpreted to reflect suboptimal nutrition (6, 40, 109). However, analysis of structural properties indicates that mechanical factors and physical activity are more important in explaining this change (see below).

CROSS-SECTIONAL GEOMETRY Because of their tubular shape, long bones (and other skeletal elements that are long relative to their width) can be modeled as hollow beams and subjected to mechanical analysis in the same way that building materials are analyzed by civil and mechanical engineers (203). In a beam (or long bone) undergoing bending or torsion (twisting), mechanical stress increases with distance from a central or neutral axis running through the shaft. Thus, the greatest mechanical stresses occurring in a lower limb bone (e.g. femur)—such as during running, walking, or lifting—are located in the outermost fibers of the bone. Therefore, a femur shaft with a more outward distribution of bone has relatively greater strength or ability to resist fracture during periods of elevated mechanical demand (179, 203).

In archaeological settings, it has long been suspected that adults having long bones with large external dimensions were physically active during life (21, 132). The analysis of cross-sectional geometric (structural) properties that measure bone strength has provided strong support for this model. On the prehistoric Georgia coast, for example, declines in external dimensions of long bones with agriculture was interpreted to reflect a decrease in mechanical demand associated with the shift from a lifeway based on hunting, gathering, and fishing to one depending at least in part on maize agriculture (132). Confirmation of this interpretation is provided by comparison of bones representing the leg (femur) and arm (humerus) whereby both sexes showed a reduction in cross-sectional properties associated with bending and torsion (139, 206). In contrast to these findings, comparisons between prehistoric foragers and agriculturalists in Alabama showed that the latter had greater bone strength (27, 29), thus indicating that the behavioral changes associated with the transition to agriculture were likely quite different in coastal (Georgia) and interior (Alabama) settings. Biomechanical analysis of human remains from other regions of North America confirm the variable nature of activity patterns in agricultural settings (33, 204).

Analysis of cross-sectional shape of long bones provides insight into the degree of mobility of past populations. Comparing human groups ranging from highly mobile hunter-gatherers to sedentary industrial populations, Ruff (201) has found that sedentary groups have a ratio approaching 1.0 in anterior-posterior bending stress (I_x) to medial-lateral bending stress (I_y) in the adult femur midshaft. Values near or at 1.0 reflect a circular shape of the bone cross-section, and represent decline in physical activities, such as running and long-distance walking. Comparisons of long-bone mechanical properties representing different subsistence technologies within specific regions indicate that agriculturalists tend to have values closer to 1.0 than do earlier hunter-gatherers (25, 33, 139). These findings suggest that, although behavioral adaptation varies between regions, human populations share a common theme involving a decline in mobility and long-distance travel in the transition from foraging to farming. Thus, the shape changes in external long-bone dimensions discussed above likely reflect an overall reduction in mobility in humans during the Holocene (203). Although the shape change occurs for human populations generally, it is especially pronounced for males. This result suggests that alterations in types of physical activity with subsistence change were greater in males than in females (201).

Comparisons of cross-sectional geometric properties and the ratio I_x/I_y for males and females show a decline in sexual dimorphism with the transition from hunting and gathering to agriculture or agriculture intensification (139, 201, 203, 204). Ethnographic documentation of mobility patterns in hunter-gatherers indicates that long-distance movement and acquisition of protein resources through hunting is nearly always done by males, whereas in agricultural and industrial societies, like females, males are involved in sedentary tasks (201, 203). Thus, the greater similarity of I_x/I_y values in males and females in agricultural populations in comparison to hunter-gatherer populations represents increasing sedentism in both sexes (203).

Structural comparison of diaphyses of left and right humeri in hunter-gatherer vs later agricultural populations within the same setting indicates a more equal distribution of mechanical demand on and use of left and right arms in later farmers. Fresia and coworkers (71) have identified a reduction in bilateral asymmetry in humeral strength in the American Southeast, a pattern that continues after European contact, when populations became more intensive maize agriculturalists. Similarly, Bridges (25, 27) documented a decrease in asymmetry of humerus strength in prehistoric Alabama, suggesting the shift to the use of both left and right arms in subsistence-related activities (e.g. pounding of maize with wooden mortars and pestles). These studies also show a decrease in sexual dimorphism in bilateral asymmetry, indicating the involvement of both women and men in similar kinds of activities in agricultural settings (27, 203, 240).

CONCLUSIONS

Within a remarkably short amount of time, agriculture became the dominant mode of subsistence, having profound effects on the landscape and populations occupying it. The shift from foraging to farming led to a reduction in health status and well-being, an increase in physiological stress, a decline in nutrition, an increase in birthrate and population growth, and an alteration of activity types and work loads. Taken as a whole, then, the popular and scholarly perception that quality of life improved with the acquisition of agriculture is incorrect.

Biological changes did not occur uniformly within populations adopting agriculture. Rather, segments of past societies were affected differentially. This pattern is especially clear in view of differences between adult females and males in dental health and activity patterns. Contrary to earlier assertions (e.g. 257), there has been a great deal of biological change in humans subsequent to the beginning of agriculture in the late Pleistocene and early Holocene. Finally, the shift to agriculture and associated biological changes in general show worldwide variability, suggesting that dietary changes and accompanying hard-tissue alterations may have been configured by highly localized circumstances.

When discussing the transition to food production, anthropology textbooks typically emphasize the social, cultural, and technological transformations that accompanied this adaptive shift. Generally not included is the biological record drawn from the study of human osteological remains. This record provides a compelling index for assessing the history of the human condition and the role that agriculture played in shaping it in recent human evolution. This review would suggest that biological changes are an essential component of lifeway transitions and should be incorporated into discussions of the consequences of the most important dietary and economic transformation in the history of humankind.

ACKNOWLEDGMENTS

I thank PS Bridges, DL Brockington, JM Calcagno, AH Goodman, DL Hutchinson, PM Lambert, CE Larsen, PW Leslie, JR Lukacs, C Meiklejohn, E Monahan, R Pastor, M Reeves, JC Rose, CB Ruff, CM Scarry, SW Simpson, VP Steponaitis, DH Ubelaker, and PL Walker for reading earlier drafts and providing valuable substantive and editorial comments. I also thank AH Goodman, PM Lambert, C Meiklejohn, T Molleson, R Pastor, JF Powell, and JC Rose for providing reprints. M Reeves and E Monahan helped locate references. Many of the ideas discussed in this article were developed during the study of North American archaeological skeletal series with support from the National Science Foundation. Collaboration with MC Griffin, DL Hutchinson,

CB Ruff, KF Russell, MJ Schoeninger, SW Simpson, MF Teaford, and DH Thomas has been instrumental in the progress of the research.

**Any Annual Review chapter, as well as any article cited in an Annual Review chapter, may be purchased from the Annual Reviews Preprints and Reprints service.
1-800-347-8007; 415-259-5017; email: arpr@class.org**

Literature Cited

1. Anderson JE. 1965. Human skeletons from Tehuacan. *Science* 148:496-97
2. Anderson JE. 1967. The human skeletons. In *The Prehistory of the Tehuacan Valley*. Vol. 1: *Environment and Subsistence*, ed. DS Byers, pp. 91-113. Austin: Univ. Texas Press
3. Angel JL. 1944. Greek teeth: ancient and modern. *Hum. Biol.* 16:283-97
4. Angel JL. 1966. Porotic hyperostosis, anemias, malaras, and marshes in the prehistoric eastern Mediterranean. *Science* 153: 760-63
5. Angel JL. 1978. Porotic hyperostosis in the eastern Mediterranean. *Med. Coll. Va. Q.* 14:10-16
6. Angel JL. 1979. Osteoarthritis in prehistoric Turkey and medieval Byzantium. In *Arthritis: Modern Concepts and Ancient Evidence*, ed. E Cockburn, H Duncan, JM Riddle. *Henry Ford Hosp. Med. J.* 27: 38-43
7. Angel JL. 1984. Health as a crucial factor in the changes from hunting to developed farming in the eastern Mediterranean. See Ref. 55, pp. 51-73
8. Armelagos GJ, Goodman AH, Jacobs KH. 1991. The origins of agriculture: population growth during a period of declining health. *Popul. Environ.* 13:9-22
9. Armelagos GJ, Rose JC. 1972. *Factors affecting tooth loss in prehistoric Nubian populations*. Presented at Annu. Meet. Soc. Am. Archaeol., Miami
10. Ashworth A, Milner PF, Waterlow JC, Walker RB. 1973. Absorption of iron from maize (*Zea mays* L.) and soya beans (*Glycine hispida* Max.) in Jamaican infants. *Br. J. Nutr.* 29:269-78
11. Beckett S, Lovell NC. 1994. Dental disease evidence for agricultural intensification in the Nubian C-Group. *Int. J. Osteoarchaeol.* 4:223-39
12. Benfer RA, Edwards DS. 1991. The principal axis method for measuring rate and amount of dental attrition: estimating juvenile or adult tooth wear from unaged adult teeth. See Ref. 121, pp. 325-40
13. Bennett KA. 1973. *The Indians of Point of Pines, Arizona: A Comparative Study of Their Physical Characteristics*. Anthropol. Pap. Univ. Ariz. 23
14. Bennike P. 1985. *Palaeopathology of Danish Skeletons*. Copenhagen: Akademisk
15. Bentley GR, Goldberg T, Jasienska G. 1993. The fertility of agricultural and non-agricultural traditional societies. *Popul. Stud.* 47:269-81
16. Bentley GR, Jasienska G, Goldberg T. 1993. Is the fertility of agriculturalists higher than that of nonagriculturalists. *Curr. Anthropol.* 34:778-85
17. Blakey ML, Armelagos GJ. 1985. Deciduous enamel defects in prehistoric Americans from Dickson Mounds: prenatal and postnatal stress. *Am. J. Phys. Anthropol.* 66:371-80
18. Bogin B. 1988. *Patterns of Human Growth*. Cambridge: Cambridge Univ. Press
19. Borgognini Tarli S, Canci A, Piperno M, Repetto E. 1993. Dati archeologici e antropologici sulle sepolture mesolitiche della Grotta dell'Uzzo (Trapani). *Bull. Paletnol. Ital.* 84:85-179
20. Boyd DCM. 1988. *A functional model for masticatory-related mandibular, dental, and craniofacial microevolutionary change derived from a selected southeastern Indian skeletal temporal series*. PhD thesis. Univ. Tenn., Knoxville
21. Boyd DCM, Boyd CC Jr. 1989. A comparison of Tennessee Archaic and Mississippian maximum femoral lengths and mid-shaft diameters: subsistence change and postcranial variability. *Southeast. Archaeol.* 8:107-16
22. Brace CL, Rosenberg KR, Hunt KD. 1987. Gradual change in human tooth size in the late Pleistocene and post Pleistocene. *Evolution* 41:705-20
23. Brace CL, Smith SL, Hunt KD. 1991. What big teeth you had Grandma! Human tooth size, past and present. See Ref. 121, pp. 33-57
24. Braidwood RJ. 1960. The agricultural revolution. *Sci. Am.* 203(3):130-48
25. Bridges PS. 1985. *Changes in long bone structure with the transition to agriculture:*

- implications for prehistoric activities. PhD thesis. Univ. Mich., Ann Arbor
26. Bridges PS. 1989. Spondylolysis and its relationship to degenerative joint disease in the prehistoric southeastern United States. *Am. J. Phys. Anthropol.* 79:321-29
 27. Bridges PS. 1989. Changes in activities with the shift to agriculture in the southeastern United States. *Curr. Anthropol.* 30: 385-94
 28. Bridges PS. 1991. Degenerative joint disease in hunter-gatherers and agriculturalists from the southeastern United States. *Am. J. Phys. Anthropol.* 85:379-91
 29. Bridges PS. 1991. Skeletal evidence of changes in subsistence activities between the Archaic and Mississippian time periods in northwestern Alabama. See Ref. 194, pp. 89-101
 30. Bridges PS. 1992. Prehistoric arthritis in the Americas. *Annu. Rev. Anthropol.* 21: 67-91
 31. Bridges PS. 1995. *Biomechanical changes in long bone diaphyses with the intensification of agriculture in the lower Illinois Valley*. Presented at Annu. Meet. Am. Assoc. Phys. Anthropol., Oakland
 32. British Association for the Advancement of Science. 1916. Is the British facial form changing? *Nature* 98:198
 33. Brock SL, Ruff CB. 1988. Diachronic patterns of change in structural properties of the femur in the prehistoric American Southwest. *Am. J. Phys. Anthropol.* 75: 113-27
 34. Brothwell DR. 1981. *Digging up Bones: The Excavation, Treatment and Study of Human Skeletal Remains*, 3e. Ithaca, NY: Cornell Univ. Press
 35. Buikstra JE, Konigsberg LW. 1985. Paleodemography: critiques and controversies. *Am. Anthropol.* 87:316-33
 36. Buikstra JE, Konigsberg LW, Bullington J. 1986. Fertility and the development of agriculture in the prehistoric Midwest. *Am. Antiq.* 51:528-46
 37. Buikstra JE, Williams S. 1991. Tuberculosis in the Americas: current perspectives. In *Human Paleopathology: Current Syntheses and Future Options*, ed. DJ Ortner, AC Aufderheide, pp. 161-72. Washington, DC: Smithsonian Inst. Press
 38. Bullington J. 1991. Deciduous dental microwear of prehistoric juveniles from the lower Illinois River Valley. *Am. J. Phys. Anthropol.* 84:59-73
 39. Burr DB, Ruff CB, Thompson DD. 1990. Patterns of skeletal histologic change through time: comparison of an archaic Native American population with modern populations. *Anat. Rec.* 226:307-13
 40. Buxton LHD. 1938. Platymelia and platycnemia. *J. Anat.* 73:31-36
 41. Calcagno JM. 1989. *Mechanisms of Human Dental Reduction: A Case Study from Post-Pleistocene Nubia*. Univ. Kans. Publ. Anthropol. 18
 42. Calcagno JM, Gibson KR. 1991. Selective compromise: evolutionary trends and mechanisms in hominid tooth size. See Ref. 121, pp. 59-76
 43. Campbell KL, Wood JW. 1988. Fertility in traditional societies. In *Natural Human Fertility: Social and Biological Determinants*, ed. P Diggory, S Teper, M Potts, pp. 39-69. London: Macmillan
 44. Carlson DS. 1976. Temporal variation in prehistoric Nubian crania. *Am. J. Phys. Anthropol.* 45:467-84
 45. Carlson DS, Armelagos GJ, Van Gerven DP. 1974. Factors influencing the etiology of cribra orbitalia in prehistoric Nubia. *J. Hum. Evol.* 3:405-10
 46. Carlson DS, Van Gerven DP. 1977. Masticatory function and post-Pleistocene evolution in Nubia. *Am. J. Phys. Anthropol.* 46: 495-506
 47. Carlson DS, Van Gerven DP. 1979. Diffusion, biological determinism, and biocultural adaptation in the Nubian Corridor. *Am. Anthropol.* 81:561-80
 48. Cassidy CM. 1984. Skeletal evidence for prehistoric subsistence adaptation in the central Ohio River valley. See Ref. 55, pp. 307-45
 49. Clark GA, Hall NR, Armelagos GJ, Borkan GA, Panjabi MM, Wetzel FT. 1986. Poor growth prior to early childhood: decreased health and life-span in the adult. *Am. J. Phys. Anthropol.* 70:145-60
 50. Clark JD, Brandt SA, eds. 1984. *From Hunters to Farmers: The Causes and Consequences of Food Production in Africa*. Berkeley: Univ. Calif. Press
 51. Clarke NG, Hirsch RS. 1991. Physiological, pulpal, and periodontal factors influencing alveolar bone. See Ref. 121, pp. 241-66
 52. Cohen MN. 1977. *The Food Crisis in Prehistory: Overpopulation and the Origins of Agriculture*. New Haven, CT: Yale Univ. Press
 53. Cohen MN. 1989. *Health and the Rise of Civilization*. New Haven, CT: Yale Univ. Press
 54. Cohen MN, Armelagos GJ. 1984. Paleopathology at the origins of agriculture: editors' summation. See Ref. 55, pp. 585-601
 55. Cohen MN, Armelagos GJ, eds. 1984. *Paleopathology at the Origins of Agriculture*. Orlando, FL: Academic
 56. Cook DC. 1984. Subsistence and health in the lower Illinois Valley: osteological evidence. See Ref. 55, pp. 237-69
 57. Corruccini RS. 1991. Anthropological aspects of orofacial and occlusal variations and anomalies. See Ref. 121, pp. 295-323
 58. Cowan CW, Watson PJ, eds. 1992. *The*

- Origins of Agriculture: An International Perspective.* Washington, DC: Smithsonian Inst. Press
59. Cybulski JS. 1977. Cribra orbitalia, a possible sign of anemia in early historic native populations of the British Columbia coast. *Am. J. Phys. Anthropol.* 47:31–40
 60. Cybulski JS. 1992. *A Greenville Burial Ground: Human Remains and Mortuary Elements in British Columbia Coast Prehistory.* Natl. Mus. Man Mercury Ser., Archaeol. Surv. Can. Pap. 146
 61. Danforth ME. 1989. *A comparison of childhood health patterns in the Late Classic and Colonial Maya using enamel microdefects.* PhD thesis. Ind. Univ., Bloomington
 62. y'Edynak G. 1978. Culture, diet, and dental reduction in Mesolithic forager-fishers of Yugoslavia. *Curr. Anthropol.* 19:616–18
 63. y'Edynak G. 1989. Yugoslav Mesolithic dental reduction. *Am. J. Phys. Anthropol.* 78:17–36
 64. y'Edynak G, Fleisch S. 1983. Microevolution and biological adaptability in the transition from food-collecting to food-producing in the Iron Gates of Yugoslavia. *J. Hum. Evol.* 12:279–96
 65. Eisenberg LE. 1991. Mississippian cultural terminations in Middle Tennessee: what the bioarchaeological evidence can tell us. See Ref. 194, pp. 70–88
 66. Eveleth PB, Tanner JM. 1990. *World-Wide Variation in Human Growth.* Cambridge: Cambridge Univ. Press. 2nd ed.
 67. Food and Agricultural Organization. 1970. *Amino-Acid Content on Foods and Biological Data on Proteins.* Rome: FAO
 68. Fogel RW, Engerman SL, Floud R, Friedman G, Margo RA, et al. 1983. Secular changes in American and British stature and nutrition. *J. Interdiscip. Hist.* 14: 445–81
 69. Frayer DW. 1978. *The Evolution of the Dentition in Upper Paleolithic and Mesolithic Europe.* Univ. Kans. Publ. Anthropol. 10
 70. Frayer DW. 1988. Caries and oral pathologies at the Mesolithic sites of Muge: Cabeço da Arruda and Moita do Sebastiao. *Trab. Antrop. Etnol.* 27:9–25
 71. Fresia AE, Ruff CB, Larsen CS. 1990. Temporal decline in bilateral asymmetry of the upper limb on the Georgia coast. In *The Archaeology of Santa Catalina de Guale. 2: Biocultural Interpretations of a Population in Transition*, ed. CS Larsen, pp. 121–32. *Anthropol. Pap. Am. Mus. Nat. Hist.* 68
 72. Frisancho AR, Garn SM, Ascoli W. 1970. Childhood retardation resulting in reduction of adult body size to lesser adolescent skeletal delay. *Am. J. Phys. Anthropol.* 33: 325–36
 73. Fritz GJ. 1990. Multiple pathways to farming in precontact eastern North America. *J. World Prehist.* 4:387–435
 74. Fritz GJ. 1994. Are the first American farmers getting younger? *Curr. Anthropol.* 35:305–9
 75. Fritz GJ. 1994. The value of archaeological plant remains for paleodietary reconstruction. See Ref. 229, pp. 21–33
 76. Garn SM. 1970. *The Earlier Gain and Later Loss of Cortical Bone.* Springfield, IL: Thomas
 77. Garn SM, Osborne RH, McCabe KD. 1979. The effect of prenatal factors on crown dimensions. *Am. J. Phys. Anthropol.* 51:665–78
 78. Garn SM, Rohmann CG, Behar M, Viteri F, Guzman MA. 1964. Compact bone deficiency in protein-calorie malnutrition. *Science* 145:1444–45
 79. Gebauer AB, Price TD, eds. 1992. *Transitions to Agriculture in Prehistory.* Madison, WI: Prehistory
 - 79a. Gilbert RI Jr, Mielke JH, eds. 1985. *The Analysis of Prehistoric Diets.* Orlando, FL: Academic
 80. Goodman AH. 1994. Cartesian reductionism and vulgar adaptationism: issues in the interpretation of nutritional status in prehistory. See Ref. 229, pp. 163–77
 81. Goodman AH, Armelagos GJ. 1988. Childhood stress, cultural buffering, and decreased longevity in a prehistoric population. *Am. Anthropol.* 90:936–44
 82. Goodman AH, Capasso LL, eds. 1992. *Recent Contributions to the Study of Enamel Developmental Defects. J. Paleopathol. Monogr. Publ.* 2
 83. Goodman AH, Lallo J, Armelagos GJ, Rose JC. 1984. Health changes at Dickson Mounds, Illinois (A.D. 950–1300). See Ref. 55, pp. 271–305
 84. Goodman AH, Martinez C, Chavez A. 1991. Nutritional supplementation and the development of linear enamel hypoplasias in children from Tezonteopan, Mexico. *Am. J. Clin. Nutr.* 53:773–81
 85. Goodman AH, Pelto GH, Allen LH, Chavez A. 1992. Socioeconomic and anthropometric correlates of linear enamel hypoplasia in children from Solis, Mexico. See Ref. 82, pp. 373–80
 86. Goodman AH, Rose JC. 1990. Assessment of systemic physiological perturbations from dental enamel hypoplasias and associated histological structures. *Yearb. Phys. Anthropol.* 33:59–110
 87. Goodman AH, Rose JC. 1991. Dental enamel hypoplasias as indicators of nutritional status. See Ref. 121, pp. 279–94
 88. Gracey M. 1987. Normal growth and nutrition. In *Nutrition and the Quality of Life*, ed. GH Bourne. *World Rev. Nutr. Diet.* 49: 160–210
 89. Greene DL. 1972. Dental anthropology of

- early Egypt and Nubia. *J. Hum. Evol.* 1: 315–24
90. Guagliardo MF. 1982. Tooth crown size differences between age groups: a possible new indicator of stress in skeletal samples. *Am. J. Phys. Anthropol.* 58:383–89
 91. Gualandi PB. 1992. Food habits and dental disease in an Iron-Age population. *Anthropol. Anz.* 50:67–82
 92. Hanihara K, Inoue N, Ito G, Kamegai T. 1981. Microevolution and tooth to denture base discrepancy in Japanese dentition. *J. Anthropol. Soc. Nippon* 89:63–70
 93. Harris DR, Hillman GC, eds. 1989. *Foraging and Farming: The Evolution of Plant Exploitation*. London: Unwin Hyman
 94. Hartnady P, Rose JC. 1991. Abnormal tooth-loss patterns among Archaic-period inhabitants of the lower Pecos region, Texas. See Ref. 121, pp. 267–78
 95. Hassan FA. 1981. *Demographic Archaeology*. New York: Academic
 96. Hildebolt CF, Molnar S. 1991. Measurement and description of periodontal disease in anthropological studies. See Ref. 121, pp. 225–40
 97. Hillson S. 1979. Diet and dental disease. *World Archaeol.* 11:147–62
 98. Hillson S. 1986. *Teeth*. Cambridge: Cambridge Univ. Press
 99. Himes JH, Martorell R, Habicht J-P, Yarbrough C, Malina RM, Klein RE. 1975. Patterns of cortical bone growth in moderately malnourished preschool children. *Hum. Biol.* 47:337–50
 100. Hinton RJ. 1981. Form and patterning of anterior tooth wear among aboriginal human groups. *Am. J. Phys. Anthropol.* 54: 555–64
 101. Hinton RJ. 1981. Temporomandibular joint size adaptations in prehistoric Tennessee Indians. *Tenn. Anthropol.* 6:89–111
 102. Hinton RJ. 1982. Differences in interproximal and occlusal tooth wear among prehistoric Tennessee Indians: implications for masticatory function. *Am. J. Phys. Anthropol.* 57:103–15
 103. Hinton RJ. 1983. Relationships between mandibular joint size and craniofacial size in human groups. *Arch. Oral Biol.* 28: 37–43
 104. Hinton RJ, Carlson DS. 1979. Temporal changes in human temporomandibular joint size and shape. *Am. J. Phys. Anthropol.* 50:325–34
 105. Hinton RJ, Smith MO, Smith FH. 1980. Tooth size changes in prehistoric Tennessee Indians. *Hum. Biol.* 52:229–45
 106. Hodges DC. 1989. *Agricultural Intensification and Prehistoric Health in the Valley of Oaxaca, Mexico*. *Mem. Mus. Anthropol. Univ. Mich.* 22
 107. Hojo T. 1989. Dietary differences and microwear on the teeth of Late Stone Age and early modern people from western Japan. *Scann. Microsc.* 3:623–28
 108. Hough AJ, Sokoloff L. 1989. Pathology of osteoarthritis. In *Arthritis and Allied Conditions*, ed. DJ McCarthy, pp. 1571–94. Philadelphia: Lea & Febiger. 11th ed.
 109. Hoyme LE, Bass WM. 1962. Human skeletal remains from the Tollifero (Ha6) and Clarksville (Mc14) sites, John H. Kerr Reservoir basin, Virginia. *Bur. Am. Ethnol. Bull.* 182:329–400
 110. Hummert JR. 1983. Skeletal growth and dietary stress among subadults from Nubia's Batn el Hajar. *Am. J. Phys. Anthropol.* 62:167–76
 111. Hummert JR, Van Gerven DP. 1983. Skeletal growth in a Medieval population from Sudanese Nubia. *Am. J. Phys. Anthropol.* 60:471–78
 112. Huss-Ashmore R, Johnston FE. 1985. Bioanthropological research in developing countries. *Annu. Rev. Anthropol.* 14: 475–528
 113. Hutchinson DL, Larsen CS. 1988. Determination of stress episode duration from linear enamel hypoplasias: a case study from St. Catherines Island, Georgia. *Hum. Biol.* 60:93–110
 114. Inoue N, Ito G, Kamegai T. 1986. Dental pathology of hunter-gatherers and early farmers in prehistoric Japan. In *Prehistoric Hunter-Gatherers in Japan: New Research Methods*, ed. T Akazawa, CM Aikens. *Univ. Tokyo Univ. Mus. Bull.* 27:163–98
 115. Jacobs K. 1994. Human dento-gnathic metric variation in Mesolithic/Neolithic Ukraine: possible evidence of demic diffusion in the Dnieper Rapids region. *Am. J. Phys. Anthropol.* 95:1–26
 116. Johansson SR, Horowitz S. 1986. Estimating mortality in skeletal populations: influence of the growth rate on the interpretation of levels and trends during the transition to agriculture. *Am. J. Phys. Anthropol.* 71: 233–50
 117. Jurmain RD. 1977. Stress and etiology of osteoarthritis. *Am. J. Phys. Anthropol.* 46: 353–66
 118. Jurmain RD. 1977. Paleoepidemiology of degenerative knee disease. *Med. Anthropol.* 1:1–14
 119. Jurmain RD. 1980. The pattern of involvement of appendicular degenerative joint disease. *Am. J. Phys. Anthropol.* 53:143–50
 120. Katz SH, Hediger ML, Valleroy LA. 1974. Traditional maize processing techniques in the New World. *Science* 184:765–73
 121. Kelley MA, Larsen CS, eds. 1991. *Advances in Dental Anthropology*. New York: Wiley-Liss
 122. Kennedy KAR. 1983. Morphological variations in ulnar supinator crests and fossae as identifying markers of occupational stress. *J. Forensic Sci.* 28:871–76

123. Kennedy KAR. 1984. Growth, nutrition, and pathology in changing paleodemographic settings in South Asia. See Ref. 55, pp. 169–92
124. Kieser JA. 1990. *Human Adult Odontometrics: The Study of Variation in Adult Tooth Size*. Cambridge: Cambridge Univ. Press
125. Klepinger LL. 1984. Nutritional assessment from bone. *Annu. Rev. Anthropol.* 13: 75–96
126. Kobayashi K. 1967. *Trend in the Length of Life Based on Human Skeletons from Prehistoric to Modern Times in Japan*. Univ. Tokyo J. Fac. Sci., Sec. V 3(2)
127. Lallo JW, Armelagos GJ, Rose JC. 1977. The role of diet, disease, and physiology in the origin of porotic hyperostosis. *Hum. Biol.* 49:471–84
128. Lallo JW, Rose JC. 1979. Patterns of stress, disease and mortality in two prehistoric populations from North America. *J. Hum. Evol.* 8:323–35
129. Lambert PM. 1993. Health in prehistoric populations of the Santa Barbara Channel Islands. *Am. Antiq.* 58:509–22
130. Lambert PM. 1994. *War and peace on the western front: a study of violent conflict and its correlates in prehistoric hunter-gatherer societies of coastal southern California*. PhD thesis. Univ. Calif., Santa Barbara
131. Lambert PM, Walker PL. 1991. Physical anthropological evidence for the evolution of social complexity in coastal southern California. *Antiquity* 65:963–73
132. Larsen CS. 1982. *The Anthropology of St. Catherine's Island. 3: Prehistoric Human Biological Adaptation*. *Anthropol. Pap. Am. Mus. Nat. Hist.* 57(3)
133. Larsen CS. 1983. Behavioural implications of temporal change in cariogenesis. *J. Archaeol. Sci.* 10:1–8
134. Larsen CS. 1983. Deciduous tooth size and subsistence change in prehistoric Georgia coast populations. *Curr. Anthropol.* 24: 225–26
135. Larsen CS. 1984. Health and disease in prehistoric Georgia: the transition to agriculture. See Ref. 55, pp. 367–92
136. Larsen CS. 1987. Bioarchaeological interpretations of subsistence economy and behavior from human skeletal remains. In *Advances in Archaeological Method and Theory*, ed. MB Schiffer, 10:339–445. San Diego: Academic
137. Larsen CS, Harn DE. 1994. Health in transition: disease and nutrition in the Georgia Bight. See Ref. 229, pp. 222–34
138. Larsen CS, Hutchinson DL. 1992. Dental evidence for physiological disruption: biocultural interpretations from the eastern Spanish Borderlands. See Ref. 82, pp. 151–69
139. Larsen CS, Ruff CB. 1994. The stresses of conquest in Spanish Florida: structural adaptation and change before and after contact. In *In the Wake of Contact: Biological Responses to Conquest*, ed. CS Larsen, GR Milner, pp. 21–34. New York: Wiley-Liss
140. Larsen CS, Schoeninger MJ, Hutchinson DL, Russell KF, Ruff CB. 1990. Beyond demographic collapse: biological adaptation and change in native populations of La Florida. In *Columbian Consequences*. Vol. 2: *Archaeological and Historical Perspectives on the Spanish Borderlands East*, ed. DH Thomas, pp. 409–28. Washington, DC: Smithsonian Inst. Press
141. Larsen CS, Shavit R, Griffin MC. 1991. Dental caries evidence for dietary change: an archaeological context. See Ref. 121, pp. 179–202
142. Lee RB. 1968. What hunters do for a living, or, how to make out on scarce resources. In *Man the Hunter*, ed. RB Lee, I DeVore, pp. 30–48. Chicago: Aldine
143. Lovejoy CO, Burstein AH, Heiple KG. 1976. The biomechanical analysis of bone strength: a method and its application to platycnemia. *Am. J. Phys. Anthropol.* 44: 489–506
144. Lubell D, Jackes M, Schwarcz H, Knyf M, Meiklejohn C. 1994. The Mesolithic-Neolithic transition in Portugal: isotopic and dental evidence of diet. *J. Archaeol. Sci.* 21:201–16
145. Lukacs JR. 1985. Tooth size variation in prehistoric India. *Am. Anthropol.* 87:1–15
146. Lukacs JR. 1990. On hunter-gatherers and their neighbors in prehistoric India: contact and pathology. *Curr. Anthropol.* 31:183–86
147. Lukacs JR. 1992. Dental paleopathology and agricultural intensification in South Asia: new evidence from Bronze Age Harappa. *Am. J. Phys. Anthropol.* 87: 133–50
148. Lukacs JR, Retief DH, Jarrige J-F. 1985. Dental disease in prehistoric Baluchistan. *Natl. Geogr. Res.* Spring:184–97
149. Lukacs JR, Schultz M, Hemphill BE. 1989. Dental pathology and dietary patterns in Iron Age northern Pakistan. In *South Asian Archaeology 1985*, ed. P Sorensen, K Frifelt, pp. 475–96. London: Curzon
150. Maat GJR, Van der Velde EA. 1987. The caries-attribution competition. *Int. J. Anthropol.* 2:281–92
151. Macchiarelli R, Bondioli L. 1986. Post-Pleistocene reductions in human dental structure: a reappraisal in terms of increasing population density. *Hum. Evol.* 1: 405–17
152. Martin DL, Armelagos GJ, King JR. 1979. Degenerative joint disease of the long bones in Dickson Mounds. In *Arthritis: Modern Concepts and Ancient Evidence*, ed. E Cockburn, H Duncan, JM Riddle. *Henry Ford Hosp. Med. J.* 27:60–63
153. Martin DL, Goodman AH, Armelagos GJ.

1985. Skeletal pathologies as indicators of quality and quantity of diet. See Ref. 79a, pp. 227–79
154. Martin RB, Burr DB. 1989. *Structure, Function, and Adaptation of Compact Bone*. New York: Raven
 155. Meiklejohn C, Baldwin JH, Schentag CT. 1988. Caries as a probable dietary marker in the western European Mesolithic. In *Diet and Subsistence: Current Archaeological Perspectives*, ed. BV Kennedy, GM LeMoine, pp. 273–79. *Proc. 19th Annu. Chacmool Conf., Univ. Calgary*
 156. Meiklejohn C, Brinch Petersen E, Alexandersen V. 1995. The later Mesolithic population of Sjaelland, Denmark, and the Neolithic transition. In *The Origins of Farming in the Baltic Zone*, ed. M Zvelebil, L Domanska, R Dennell. Oxford: Oxbow. In press
 157. Meiklejohn C, Schentag C, Venema A, Key P. 1984. Socioeconomic change and patterns of pathology and variation in the Mesolithic and Neolithic of western Europe: some suggestions. See Ref. 55, pp. 75–100
 158. Meiklejohn C, Wyman JM, Jacobs K, Jackes MK. 1995. Issues in the archeological demography of the agricultural transition in western and northern Europe: a view from the Mesolithic. In *Integrating Archaeological Demography: Multidisciplinary Approaches to Prehistoric Populations*, ed. R Paine. *Cent. Archaeol. Invest., S. Ill. Univ., Occas. Pap.* 24. In press
 159. Meiklejohn C, Wyman JM, Schentag CT. 1992. Caries and attrition: dependent or independent variables? *Int. J. Anthropol.* 7:17–22
 160. Meiklejohn C, Zvelebil M. 1991. Health status of European populations at the agricultural transition and the implications for the adoption of farming. In *Health in Past Societies*, ed. H Bush, M Zvelebil, pp. 129–45. *BAR Int. Ser.* 567
 161. Mensforth RP, Lovejoy CO, Lallo JW, Armelagos GJ. 1978. The role of constitutional factors, diet, and infectious disease in the etiology of porotic hyperostosis and periosteal reactions in prehistoric infants and children. *Med. Anthropol.* 2: 1–59
 162. Merbs CF. 1983. *Patterns of Activity-Induced Pathology in a Canadian Inuit Population*. *Natl. Mus. Man Mercury Ser., Archaeol. Surv. Can. Pap.* 119
 163. Merbs CF. 1989. Spondylolysis: its nature and anthropological significance. *Int. J. Anthropol.* 4:163–69
 164. Miller RJ. 1985. Lateral epicondylitis in a prehistoric central Arizona Indian population from Nuvakwewtaqa (Chavez Pass). In *Health and Disease in the Prehistoric Southwest*, ed. CF Merbs, RJ Miller, pp. 391–400. *Ariz. State Univ. Anthropol. Res. Pap.* 4
 165. Milner GR. 1984. Dental caries in the permanent dentition of a Mississippian period population from the American Midwest. *Coll. Anthropol.* 8:77–91
 166. Milner GR. 1991. Health and cultural change in the late prehistoric American Bottom, Illinois. See Ref. 194, pp. 52–69
 167. Milner GJ, Anderson E, Smith VG. 1991. Warfare in late prehistoric west-central Illinois. *Am. Antiq.* 56:581–603
 168. Milner GR, Humpf DA, Harpending HC. 1989. Pattern matching of age-at-death distributions in paleodemographic analysis. *Am. J. Phys. Anthropol.* 80:49–58
 169. Milner GR, Larsen CS. 1991. Teeth as artifacts of human behavior: intentional mutilation and accidental modification. See Ref. 121, pp. 357–78
 170. Molleson T. 1989. Seed preparation in the Mesolithic: the osteological evidence. *Antiquity* 63:356–62
 171. Molleson T, Jones K. 1991. Dental evidence for dietary change at Abu Hureyra. *J. Archaeol. Sci.* 18:525–39
 172. Molleson T, Jones K, Jones S. 1993. Dietary change and the effects of food preparation on microwear patterns in the Late Neolithic of Abu Hureyra, northern Syria. *J. Hum. Evol.* 24:455–68
 173. Molnar S. 1972. Tooth wear and culture: a survey of tooth functions among some prehistoric populations. *Curr. Anthropol.* 13: 511–26
 174. Molnar S, Molnar I. 1985. Observations of dental diseases among prehistoric populations of Hungary. *Am. J. Phys. Anthropol.* 67:51–63
 175. Moorrees CFA. 1957. *The Aleut Dentition: A Correlative Study of Dental Characteristics in an Eskimoid People*. Cambridge, MA: Harvard Univ. Press
 176. Newbrun E. 1982. Sugar and dental caries: a review of human studies. *Science* 217: 418–23
 177. Newman MT, Snow CE. 1942. Preliminary report on the skeletal material from Pickwick Basin, Alabama. In *An Archeological Survey of Pickwick Basin in the Adjacent Portions of the States of Alabama, Mississippi and Tennessee*, ed. WS Webb, DL DeJarnette, pp. 393–507. *Bur. Am. Ethnol. Bull.* 129
 178. Nickens PR. 1976. Stature reduction as an adaptive response to food production in Mesoamerica. *J. Archaeol. Sci.* 3:31–41
 179. Nordin M, Frankel VH. 1980. Biomechanics of whole bones and bone tissue. In *Basic Biomechanics of the Skeletal System*, ed. VH Frankel, M Nordin, pp. 15–60. Philadelphia: Lea & Febiger
 180. Norr L. 1984. Prehistoric subsistence and health status of coastal peoples from the

- Panamanian isthmus of lower Central America. See Ref. 55, pp. 463–90
181. Ortner DJ. 1968. Description and classification of degenerative bone changes in the distal joint surfaces of the humerus. *Am. J. Phys. Anthropol.* 28:139–56
 182. Ortner DJ. 1992. Skeletal paleopathology: probabilities, possibilities, and impossibilities. In *Disease and Demography in the Americas*, ed. JW Verano, DH Ubelaker, pp. 5–13. Washington, DC: Smithsonian Inst. Press
 183. Owsley DW. 1991. Temporal variation in femoral cortical thickness of North American Plains Indians. See Ref. 37, pp. 105–10
 184. Pastor R. 1992. Dietary adaptations and dental microwear in Mesolithic and Chalcolithic South Asia. In *Culture, Ecology and Dental Anthropology*, ed. JR Lukacs. *J. Hum. Ecol.* (Special Issue) 2:215–28
 185. Pastor R. 1993. *Dental microwear among prehistoric inhabitants of the Indian subcontinent: a quantitative and comparative analysis*. PhD thesis. Univ. Ore., Eugene
 186. Pastor R. 1994. *A multivariate dental microwear analysis of prehistoric groups from the Indian subcontinent*. Presented at Annu. Meet. Am. Assoc. Phys. Anthropol., Denver
 187. Patterson DK Jr. 1984. *A Diachronic Study of Dental Palaeopathology and Attritional Status of Prehistoric Ontario Pre-Iroquois and Iroquois Populations*. Natl. Mus. Man Mercury Ser., Archaeol. Surv. Can. Pap. 122
 188. Perzigian AJ, Tench PA, Braun DJ. 1984. Prehistoric health in the Ohio River valley. See Ref. 55, pp. 347–66
 189. Pickering RB. 1984. *Patterns of degenerative joint disease in Middle Woodland, Late Woodland, and Mississippian skeletal series from the Lower Illinois Valley*. PhD thesis. Northwestern Univ., Evanston, Ill.
 190. Pierce LKC. 1987. *A comparison of the pattern of involvement of degenerative joint disease between an agricultural and non-agricultural skeletal series*. PhD thesis. Univ. Tenn., Knoxville
 191. Powell JF. 1995. Bioarchaeological analyses of human skeletal remains from the Mitchell Ridge site. In *Aboriginal Life and Culture on the Upper Texas Coast: Archaeology of the Mitchell Ridge Site, 41GV66, Galveston Island*, ed. RA Ricklis, JF Powell, BW Baker, JA Huebner. Corpus Christi, TX: Coastal Archaeol. Res.
 192. Powell ML. 1985. The analysis of dental wear and caries for dietary reconstruction. See Ref. 79a, pp. 307–38
 193. Powell ML. 1988. *Status and Health in Prehistory: A Case Study of the Moundville Chiefdom*. Washington, DC: Smithsonian Inst. Press
 194. Powell ML, Bridges PS, Mires AMW, eds. 1991. *What Mean These Bones? Studies in Southeastern Bioarchaeology*. Tuscaloosa: Univ. Ala. Press
 195. Rathbun TA. 1984. Skeletal pathology from the Paleolithic through the Metal Ages in Iran and Iraq. See Ref. 55, pp. 137–67
 196. Reinhard KJ. 1992. Patterns of diet, parasitism, and anemia in prehistoric west North America. In *Diet, Demography, and Disease: Changing Perspectives on Anemia*, ed. P Stuart-Macadam, S Kent, pp. 219–58. Hawthorne, NY: Aldine
 197. Rose JC, Armelagos GJ, Lallo JW. 1978. Histological enamel indicator of childhood stress in prehistoric skeletal samples. *Am. J. Phys. Anthropol.* 49:511–16
 198. Rose JC, Burnett BA, Blaeuer MW, Nassaney MS. 1984. Paleopathology and the origins of maize agriculture in the Lower Mississippi Valley and Caddoan culture areas. See Ref. 55, pp. 393–424
 199. Rose JC, Condon KW, Goodman AH. 1985. Diet and dentition: developmental disturbances. See Ref. 79a, pp. 281–305
 200. Rose JC, Marks MK, Tieszen LL. 1991. Bioarchaeology and subsistence in the central and lower portions of the Mississippi Valley. See Ref. 194, pp. 7–21
 201. Ruff CB. 1987. Sexual dimorphism in human lower limb bone structure: relationship to subsistence strategy and sexual division of labor. *J. Hum. Evol.* 16:396–416
 202. Ruff CB. 1991. *Aging and Osteoporosis in Native Americans from Pecos Pueblo, New Mexico*. New York: Garland
 203. Ruff CB. 1992. Biomechanical analyses of archaeological human material. In *Skeletal Biology of Past People*, ed. SR Saunders, MA Katzenberg, pp. 41–62. New York: Wiley-Liss
 204. Ruff CB. 1994. Biomechanical analysis of northern and southern Plains femora: behavioral implications. In *Skeletal Biology in the Great Plains: Migration, Warfare, Health, and Subsistence*, ed. DW Owsley, RL Jantz, pp. 235–45. Washington, DC: Smithsonian Inst. Press
 205. Ruff CB, Hayes WC. 1982. Subperiosteal expansion and cortical remodeling of the human femur and tibia with aging. *Science* 217:945–48
 206. Ruff CB, Larsen CS, Hayes WC. 1984. Structural changes in the femur with the transition to agriculture on the Georgia coast. *Am. J. Phys. Anthropol.* 64:125–36
 207. Ruff CB, Trinkaus E, Walker A, Larsen CS. 1993. Postcranial robusticity in *Homo*. I: Temporal trends and mechanical interpretation. *Am. J. Phys. Anthropol.* 91:21–53
 208. Sandford MK, ed. 1993. *Investigations of Ancient Human Tissue: Chemical Analyses in Anthropology*. Langhorne: Gordon & Breach
 209. Sattenspiel L, Harpending H. 1983. Stable

- populations and skeletal age. *Am. Antiquity* 48:489-98
210. Saunders SR. 1992. Subadult skeletons and growth related studies. See Ref. 203, pp. 1-20
 211. Saunders SR, Hoppa RD. 1993. Growth deficit in survivors and non-survivors: biological mortality bias in subadult skeletal samples. *Yearb. Phys. Anthropol.* 36: 127-51
 212. Scarry CM, ed. 1993. *Foraging and Farming in the Eastern Woodlands*. Gainesville: Univ. Fla. Press
 213. Schmucker BJ. 1985. Dental attrition: a correlative study of dietary and subsistence patterns. In *Health and Disease in the Prehistoric Southwest*, ed. CF Merbs, RJ Miller, pp. 275-323. *Ariz. State Univ. Anthropol. Res. Pap.* 34
 214. Schwarcz HP, Schoeninger MJ. 1991. Stable isotope analyses in human nutritional ecology. *Yearb. Phys. Anthropol.* 34: 283-321
 215. Sciulli PW, Carlisle R. 1977. Analysis of the dentition from three western Pennsylvania Late Woodland sites. II. Wear and pathology. *Penn. Archaeol.* 47: 53-59
 216. Scott EC. 1979. Increase in tooth size in prehistoric coastal Peru, 10,000 B.P.-1,000 B.P. *Am. J. Phys. Anthropol.* 50:251-58
 217. Scott GR, Turner CG II. 1988. Dental anthropology. *Annu. Rev. Anthropol.* 17: 99-126
 218. Scrimshaw NS, Taylor CE, Gordon JE. 1968. *Interaction of Nutrition and Infection*. *World Health Organ. Monogr.* 57
 219. Sealy JC, Patrick MK, Morris AG, Alder D. 1992. Diet and dental caries among Later Stone Age inhabitants of the Cape Province, South Africa. *Am. J. Phys. Anthropol.* 88:123-34
 220. Simpson SW, Hutchinson DL, Larsen CS. 1990. Coping with stress: tooth size, dental defects, and age-at-death. In *The Archaeology of Mission Santa Catalina de Guale. 2: Biocultural Interpretations of a Population in Transition*, ed. CS Larsen, pp. 66-77. *Anthropol. Pap. Am. Mus. Nat. Hist.* 68
 221. Smith BD. 1989. Origins of agriculture in eastern North America. *Science* 246: 1566-71
 222. Smith BD. 1992. Hopewellian farmers of eastern North America. In *Rivers of Change: Essays on Early Agriculture in Eastern North America*, ed. BD Smith, pp. 201-48. Washington, DC: Smithsonian Inst. Press
 223. Smith BH. 1984. Patterns of molar wear in hunter-gatherers and agriculturalists. *Am. J. Phys. Anthropol.* 63:39-56
 224. Smith FH, Smith MO, Hinton RJ. 1980. Evolution of tooth size in the prehistoric inhabitants of the Tennessee Valley. In *The Skeletal Biology of Aboriginal Populations in the Southeastern United States*, ed. P Willey, FH Smith, pp. 81-103. *Tenn. Anthropol. Assoc. Misc. Pap.* 5
 225. Smith MO. 1982. *Patterns of association between oral health status and subsistence: a study of aboriginal skeletal populations from the Tennessee Valley area*. PhD thesis, Univ. Tenn., Knoxville
 226. Smith P. 1972. Diet and attrition in the Natufians. *Am. J. Phys. Anthropol.* 37: 233-38
 227. Smith P, Bar-Yosef O, Sillen A. 1984. Archaeological and skeletal evidence for dietary change during the late Pleistocene/early Holocene in the Levant. See Ref. 55, pp. 101-36
 228. Sobolik KD. 1994. Paleonutrition of the Lower Pecos region of the Chihuahuan Desert. See Ref. 229, pp. 247-64
 229. Sobolik KD, ed. 1994. *Paleonutrition: The Diet and Health of Prehistoric Americans*. *Cent. Archaeol. Invest., S. Ill. Univ., Occas. Pap.* 22
 230. Stahl AB. 1989. Plant-food processing: implications for dietary quality. In *Foraging and Farming: The Evolution of Plant Exploitation*, ed. DR Harris, GC Hillman, pp. 171-94. London: Unwin Hyman
 231. Steckel RH. 1987. Growth depression and recovery: the remarkable case of American slaves. *Ann. Hum. Biol.* 14:111-32
 232. Steegmann AT Jr. 1985. 18th century British military stature: growth cessation, selective recruiting, secular trends, nutrition at birth, cold and occupation. *Hum. Biol.* 57:77-95
 233. Stewart TD. 1969. The effects of pathology on skeletal populations. *Am. J. Phys. Anthropol.* 30:443-50
 234. Stout SD. 1983. The application of histomorphometric analysis to ancient skeletal remains. *Anthropos* 10:60-71
 235. Stout SD. 1989. Histomorphometric analysis of human skeletal remains. In *Reconstruction of Life from the Skeleton*, ed. MY Iscan, KAR Kennedy, pp. 41-52. New York: Liss
 236. Stuart-Macadam P. 1992. Anemia in past human populations. See Ref. 196, pp. 151-70
 237. Suzuki H. 1969. Microevolutional changes in the Japanese population from the prehistoric age to the present-day. *Univ. Tokyo J. Fac. Sci., Sec. V* 3:279-308
 238. Teaford MF. 1991. Dental microwear: What can it tell us about diet and dental function? See Ref. 121, pp. 341-56
 239. Townsend GC, Brown T. 1978. Heritability of permanent tooth size. *Am. J. Phys. Anthropol.* 49:497-504
 240. Trinkaus E, Churchill SE, Ruff CB. 1994. Postcranial robusticity in *Homo*: II. hu-

- meral bilateral asymmetry and bone plasticity. *Am. J. Phys. Anthropol.* 93:1–34
241. Truswell AS, Hansen JDL. 1976. Medical research among the !Kung. In *Kalahari Hunter-Gatherers*, ed. RB Lee, I DeVore, pp. 166–94. Cambridge, MA: Harvard Univ. Press
242. Turner CG II. 1979. Dental anthropological indications of agriculture among Jomon people of central Japan. *Am. J. Phys. Anthropol.* 51:619–36
243. Turner CG II. 1982. *Oral health and other biological relationships with economic systems in eastern Asian skeletal populations*. Presented at Conf. Paleopathol. Socioecon. Change at the Origins of Agric., Plattsburgh, NY
244. Turner CG II, Cadien JD. 1969. Dental chipping in Aleuts, Eskimos and Indians. *Am. J. Phys. Anthropol.* 31:303–10
245. Ubelaker DH. 1979. Skeletal evidence for kneeling in prehistoric Ecuador. *Am. J. Phys. Anthropol.* 51:679–86
246. Ubelaker DH. 1984. Prehistoric human biology of Ecuador: possible temporal trends and cultural correlations. See Ref. 55, pp. 491–513
247. Ubelaker DH. 1992. Enamel hypoplasia in Ecuador. See Ref. 82, pp. 207–17
248. Ubelaker DH. 1994. The biological impact of European contact in Ecuador. See Ref. 139, pp. 147–60
249. Van Gerven DP. 1982. The contribution of time and local geography to craniofacial variation in Nubia's Batn el Hajar. *Am. J. Phys. Anthropol.* 59:307–16
250. Van Gerven DP, Armelagos GJ, Rohr A. 1977. Continuity and change in cranial morphology of three Nubian archaeological populations. *Man* 122:270–77
251. Walker PL. 1978. A quantitative analysis of dental attrition rates in the Santa Barbara Channel area. *Am. J. Phys. Anthropol.* 48:101–6
252. Walker PL. 1986. Porotic hyperostosis in a marine-dependent California Indian population. *Am. J. Phys. Anthropol.* 69:345–54
253. Walker PL, Dean G, Shapiro P. 1991. Estimating age from tooth wear in archaeological populations. See Ref. 121, pp. 169–78
254. Walker PL, Erlandson J. 1986. Dental evidence for prehistoric dietary change on the northern Channel Islands. *Am. Antiq.* 51: 375–83
255. Walker PL, Hewlett BS. 1990. Dental health diet and social status among Central African foragers and farmers. *Am. Anthropol.* 92:383–98
256. Walker PL, Johnson J, Lambert P. 1988. Age and sex biases in the preservation of human skeletal remains. *Am. J. Phys. Anthropol.* 76:183–88
257. Washburn SL, Lancaster CS. 1966. The evolution of hunting. In *Man the Hunter*, ed. RB Lee, I DeVore, pp. 293–303. Chicago: Aldine
258. Welinder S. 1979. *Prehistoric Demography*. *Acta Arch. Lundensia* (IN8 Minore 8). Bonn: Rudolf Habelt
259. Whittington SL. 1992. Enamel hypoplasia in the low status Maya population of prehispanic Copan, Honduras. See Ref. 82, pp. 185–205
260. Willey P. 1990. *Prehistoric Warfare on the Great Plains: Skeletal Analysis of the Crow Creek Massacre Victims*. New York: Garland
261. Wolf G. 1980. Vitamin A. In *Human Nutrition*, ed. RB Alfin-Slater, D Kritchevsky, pp. 97–203. New York: Plenum
262. Wood BF. 1971. Malocclusion in the modern Alaskan Eskimo. *Am. J. Orthod.* 60: 344–54
263. Wood JW, Milner GR, Harpending HC, Weiss KM. 1992. The osteological paradox: problems of inferring prehistoric health from skeletal samples. *Curr. Anthropol.* 33:343–58
264. Wu X, Zhang Z. 1985. *Homo sapiens* remains from late Palaeolithic and Neolithic China. In *Palaeoanthropology and Palaeolithic Archaeology in the People's Republic of China*, ed. R Wu, JW Olsen, pp. 107–33. Orlando, FL: Academic
265. Yagi T, Takebe Y, Itoh M. 1989. Secular trends in physique and physical fitness in Japanese students during the last 20 years. *Am. J. Hum. Biol.* 1:581–87
266. Yamamoto M. 1992. Secular trends of enamel hypoplasia in Japanese from the prehistoric to modern period. See Ref. 82, pp. 231–38